

Radiation-induced solitary waves in hot plasmas

F.V.Prigara

Institute of Microelectronics and Informatics, Russian Academy of Sciences,

*21 Universitetskaya, 150007 Yaroslavl, Russia**

(Dated: February 2, 2008)

Abstract

It is argued that, in a hot plasma interacting with thermal radiation, there exist the radiation-induced solitary waves different from the types of nonlinear waves in plasmas known so far. There are extensive observational evidence for the propagation of these specific density waves in the hot plasmas of various astrophysical objects. Radiation produces also non-adiabatic density perturbations analogues to the second sound in the liquid helium.

PACS numbers: 52.25.Qs, 52.35.Sb, 95.30.Gv, 95.30.Jx.

*Electronic address: fprigara@imras.yar.ru

In hot plasmas, the interaction between radiation and matter plays a very important role. In fact, a hot plasma is a four-component medium which includes the electron and ion fluids, magnetic field, and radiation. Each of the first three components produces its own branch of linear and nonlinear waves in plasmas [1,2]. Plasma waves and Langmuir envelope solitons correspond to the electron fluid. Ion sound and ion-sound shocks correspond to the ion fluid. Magnetic field is responsible for magnetosonic and Alfvén waves, and also for the magnetosonic shocks and solitary waves [3].

Quite similarly, the radiation component of a hot plasma produces its own branches of linear and nonlinear waves. Linear radiation-induced waves are non-adiabatic density perturbations, or entropy density waves, similar to the second sound in a superfluid [4]. Here the density of entropy can be interpreted as the number of photons per one nucleon so as in the Friedmann's cosmological models with hot matter [5,6].

The nonlinear branch includes radiation-induced density waves in hot plasmas which are considered in more detail below.

To introduce the radiation-induced density waves in hot plasmas, consider at first the energy distribution of atoms (ions) in the field of thermal radiation. We restrict ourselves to the case of the sufficiently rarefied plasma when the line broadening due to the gas pressure is negligible.

Recently, the energy distribution of atoms in the field of thermal blackbody radiation was obtained [7] in the form

$$N/N_0 = \sigma_a \omega^2 / (2\pi c^2) (\exp(\hbar\omega/kT) - 1), \quad (1)$$

where N_0 is the population of the ground state E_0 , N is the population of the energy level $E = E_0 + \hbar\omega$, σ_a is the absorption cross-section, $\hbar$ is the Planck constant, and T is the radiation temperature.

This distribution is valid in the range $\hbar\omega/kT \geq 1$, since in the limit $\hbar\omega/kT \rightarrow 0$ the line width is going to infinity, that indicates the violation of the one-particle approximation used in [7].

The function (1) has a maximum at $\hbar\omega_m = 1.6kT$. When the temperature exceeds the critical value of $T_0 = 3 \times 10^7 K$ (the inversion temperature), the population of the energy level E exceeds the population of the ground state E_0 . Since the function (1) is increasing in the range $\omega < \omega_m$, the inversion of the energy level population is produced also in

some vicinity of ω_m (below ω_m). This suggests the maser amplification of thermal radio emission in continuum by a hot plasma with the temperature exceeding the critical value T_0 . Maser amplification in compact radio sources was assumed earlier in [8] based on the high brightness temperatures of active galactic nuclei. Since a hot plasma in an accretion disk is concentrated nearby the central energy source, maser amplification is characteristic for compact radio sources.

It is clear that, when the temperature of a plasma is below T_0 , the radio flux is very small, and when the temperature exceeds T_0 , radio emission is on. This an on-off cycle is detected in the radio pulsar PSR B1259-63 [9]. Similar is an on-off cycle in X-ray pulsars, e.g., the 35-day cycle in Her X-1. It implies that X-ray emission from X-ray pulsars is produced by the laser amplification in continuum which is quite analogues to maser amplification at radio wavelengths.

It was shown recently [7] that thermal emission has a stimulated character. According to this conception thermal emission from non-uniform gas is produced by an ensemble of individual emitters. Each of these emitters is an elementary resonator the size of which has an order of magnitude of mean free path l of photons

$$l = \frac{1}{n\sigma} \quad (2)$$

where n is the number density of particles and σ is the absorption cross-section.

The emission of each elementary resonator is coherent, with the wavelength

$$\lambda = l, \quad (3)$$

and thermal emission of gaseous layer is incoherent sum of radiation produced by individual emitters.

An elementary resonator emits in the direction opposite to the direction of the density gradient. The wall of the resonator corresponding to the lower density is half-transparent due to the decrease of absorption with the decreasing gas density.

The condition (3) implies that the radiation with the wavelength λ is produced by the gaseous layer with the definite number density of particles n .

The condition (3) is consistent with the experimental results by Looney and Brown on the excitation of plasma waves by electron beam (see [1,10]). The wavelength of standing wave

with the Langmuir frequency of oscillations depends on the density as predicted by equation (2). The discrete spectrum of oscillations is produced by the non-uniformity of plasma and the readjustment of the wavelength to the length of resonator. From the results of experiment by Looney and Brown the absorption cross-section for plasma can be evaluated.

The product of the wavelength by density is weakly increasing with the increase of density. This may imply the weak dependence of the size of elementary resonator in terms of the wavelength upon the density or, equivalently, wavelength.

If the temperature of plasma exceeds the inversion temperature, T_0 , then the inversion of energy level population is created. In this case thermal radiation from an elementary resonator is amplified by the laser mechanism and emitted in the direction of decreasing gas density, as earlier.

The time of life for an elementary resonator has an order of magnitude

$$\tau \cong l/v_{iT} \cong l/c_s, \quad (4)$$

where v_{iT} is the ion thermal velocity, and c_s is the speed of sound. Such is the time duration of the light pulse produced by an elementary resonator.

The absorption of this radiation pulse by the nearby gaseous layer leads to the heating of a plasma and its density perturbations. The radiation pulse is then re-emitted by the new virtual elementary resonator in a plasma. In such a manner, the radiation-induced density wave propagates through the hot plasma. The width of the density wave has an order of magnitude

$$\Delta \cong c\tau \cong cl/c_s, \quad (5)$$

where c is the speed of light.

Various astrophysical objects, such as active galactic nuclei, X-ray binaries, young pulsars, have hot accretion disks (e.g., [11]). Since the density gradient in an accretion disk is directed along the radius, the radiation-induced density wave in a hot plasma of an accretion disk normally propagates as the radial density wave, though the transverse density wave is also possible and seems to be observed in some pulsars.

The inversion of energy level population associated with the radial density wave produces a moving pulse of coherent radiation with the changing wavelength determined by equations

(2) and (3). Such are the radio pulses from pulsars [12].

In fact, the inversion of energy level population corresponding to the radio band (in some vicinity of ω_m below it - see Sec. 2) exists also at the temperatures below the inversion temperature, T_0 . This conclusion is confirmed by the temperature profile in active galactic nuclei [13] which is virial, i.e.

$$T \propto r^{-1}, \quad (6)$$

where r is the distance from the central energy source. Although the temperature is decreasing with the increase of the radius, maser amplification of radio emission producing the high brightness temperatures is observed in the wide range of wavelengths corresponding, in accordance with equation (3), to the wide range of the radius scales.

However, there are no observed pulsed emission from the active galactic nuclei, contrary to the pulsars. Since the temperature profile in pulsars is different from those given by equation (6) and has a form [12]

$$dT/dr \geq 0, \quad (7)$$

it is clear, that a hot plasma is required for triggering of radio and X-ray pulses. Another feature, by which pulsars differ from active galactic nuclei, is the high ratio of the gas pressure to magnetic pressure (beta) in pulsars' plasmas [12].

It suggests that the strong magnetic field is not obligatory for the propagation of the radiation-induced density waves in hot plasmas. However, if the strong magnetic field is present, it determines the direction in which the density wave is propagating. Such seems to be the case of solar microwave bursts. The radiation-induced density wave propagates along a loop causing the strong correlation in the modulation of radio and X-ray emission over a large distance ($10^{10}cm$) in the solar corona [14].

Some X-ray binaries, e.g. Sco X-1, show weak emission lines with the variable intensities and radial velocities. These features are characteristic for non-saturated lasers. The wavelength of generated mode is determined by the size of an elementary resonator which is depending on the density. The radial density wave travelling along the radius changes the density which results in the variations of the wavelength. Convection or advection in the gaseous disk produces a two-temperature plasma in which it is possible to create the

inversion of the energy level population. This gives rise to the weak lasers similar to the weak molecular masers.

The radial density wave produces also the delay of flares at low frequencies and high frequency quasi-periodic oscillations observed in X-ray binaries.

To summarize, in a hot plasma interacting with thermal radiation, the radiation-induced solitary waves can propagate in the direction of the decreasing plasma density. In the presence of the strong magnetic field the last determines the direction of propagation for the density waves. There is extensive observational evidence for the density waves in hot plasmas of accretion disks and the solar corona. The question of detecting the radiation-induced solitary waves in the hot laboratory plasmas, e.g. produced by the intense laser beam interacting with a solid surface, remains open. An effect indicative of the radiation-induced density wave in a hot plasma is a frequency-shifting pulse of coherent radiation.

-
- [1] F.F.Chen, *Introduction to Plasma Physics and Controlled Fusion, Vol. 1: Plasma Physics* (Plenum Press, New York, 1984).
 - [2] B.B.Kadomtsev, *Collective Phenomena in Plasmas* (Nauka, Moscow, 1988).
 - [3] P.K.Shukla, B.Eliasson, M.Marklund, *Nonlinear model for magnetosonic shocklets in plasmas*, Phys. Plasmas (submitted), E-print archives, physics/0402035.
 - [4] L.D.Landau and E.M.Lifshitz, *Fluids Mechanics* (Butterworth-Heinemann, New York, 1987).
 - [5] L.B.Okun, *Leptons and Quarks* (Elsevier Science, Amsterdam, 1987).
 - [6] S.Weinberg, *Gravitation and Cosmology* (Wiley, New York, 1972).
 - [7] F.V.Prigara, in *Plasmas in the Laboratory and in the Universe*, September 16-19, 2003, Como, Italy, E-prints archives, astro-ph/0311532 (2003).
 - [8] F.V.Prigara, Astron. Nachr., **324**, No. S1, 425 (2003).
 - [9] G.J.Qiao, X.Q.Xue, R.X.Xu, H.G.Wang, and B.W.Xiao, Astron. Astrophys., **407**, L25 (2003).
 - [10] B.V.Alexeev, Usp. Fiz. Nauk, **173**, 145 (2003), Physics-Uspekhi, **46** (2003).
 - [11] F.Yuan, and A.Zdziarski, *Luminous hot accretion flows: the origin of X-ray emission of Seyfert galaxies and black hole binaries*, Month. Not. R. Astron. Soc. (submitted), E-print archives, astro-ph/0401058 (2004).

- [12] F.V.Prigara, in *Gravitation, Cosmology and Relativistic Astrophysics*, June 23-27, 2003, Kharkov, Ukraine, E-print archives, astro-ph/0307288 (2003).
- [13] N.M.Nagar, A.S.Wilson, and H.Falcke, *Astrophys. J.*, **559**, L87 (2001).
- [14] V.V.Grechnev, S.M.White, and M.R.Kundu, *Astrophys. J.*, **588**, 1163 (2003).